



FERAHTIA N.¹, ALLAOUI S.E.²

A GENERALIZATION OF A LOCALIZATION PROPERTY OF BESOV SPACES

The notion of a localization property of Besov spaces is introduced by G. Bourdaud, where he has provided that the Besov spaces $B_{p,q}^s(\mathbb{R}^n)$, with $s \in \mathbb{R}$ and $p, q \in [1, +\infty]$ such that $p \neq q$, are not localizable in the ℓ^p norm. Further, he has provided that the Besov spaces $B_{p,q}^s$ are embedded into localized Besov spaces $(B_{p,q}^s)_{\ell^p}$ (i.e., $B_{p,q}^s \hookrightarrow (B_{p,q}^s)_{\ell^p}$, for $p \geq q$). Also, he has provided that the localized Besov spaces $(B_{p,q}^s)_{\ell^p}$ are embedded into the Besov spaces $B_{p,q}^s$ (i.e., $(B_{p,q}^s)_{\ell^p} \hookrightarrow B_{p,q}^s$, for $p \leq q$). In particular, $B_{p,p}^s$ is localizable in the ℓ^p norm, where ℓ^p is the space of sequences $(a_k)_k$ such that $\|(a_k)\|_{\ell^p} < \infty$. In this paper, we generalize the Bourdaud theorem of a localization property of Besov spaces $B_{p,q}^s(\mathbb{R}^n)$ on the ℓ^r space, where $r \in [1, +\infty]$. More precisely, we show that any Besov space $B_{p,q}^s$ is embedded into the localized Besov space $(B_{p,q}^s)_{\ell^r}$ (i.e., $B_{p,q}^s \hookrightarrow (B_{p,q}^s)_{\ell^r}$, for $r \geq \max(p, q)$). Also we show that any localized Besov space $(B_{p,q}^s)_{\ell^r}$ is embedded into the Besov space $B_{p,q}^s$ (i.e., $(B_{p,q}^s)_{\ell^r} \hookrightarrow B_{p,q}^s$, for $r \leq \min(p, q)$). Finally, we show that the Lizorkin-Triebel spaces $F_{p,q}^s(\mathbb{R}^n)$, where $s \in \mathbb{R}$ and $p, q \in [1, +\infty]$ are localizable in the ℓ^p norm (i.e., $F_{p,q}^s = (F_{p,q}^s)_{\ell^p}$)

Key words and phrases: Besov spaces, Lizorkin-Triebel spaces, Localization property.

¹ Laboratory of Pures and Applied Mathematics, Department of Mathematics, Mohamed Boudiaf University of Msila, P.O. Box 166 Ichbilia, Msila 28000, Algeria

² Department of Mathematics and Informatics, Laghouat University, Laghouat 03000, Algeria
E-mail: na.ferahtia@gmail.com, shallaoui@yahoo.fr

INTRODUCTION

Functional calculus is one of the basic theory in functional analysis [5]. It has enabled to study function-analytic in topological (in particular, normed) spaces of functions. For instance, several authors such as Peetre [7], Dahlberg [4], Marcus and Mizel [6] have studied functional calculus in certain Sobolev and Besov spaces. In particular, Bourdaud [1, 2] have established a way of functional calculus in localized Besov spaces. More precisely, in [1] he has proved the following result.

Theorem 1. *Let $p, q \in [1, +\infty]$, $s \in \mathbb{R}$, and $B_{p,q}^s$ and $(B_{p,q}^s)_{\ell^p}$ are respectively the Besov and localized Besov spaces. Then*

(i) $B_{p,q}^s \hookrightarrow (B_{p,q}^s)_{\ell^p}$, for $p \geq q$,

(ii) $(B_{p,q}^s)_{\ell^p} \hookrightarrow B_{p,q}^s$, for $p \leq q$.

In particular, $B_{p,p}^s$ is localizable in the ℓ^p norm, where ℓ^p is the space of sequences $(a_k)_k$ such that $\|(a_k)\|_{\ell^p} = (\sum_{k=0}^{\infty} |a_k|^p)^{\frac{1}{p}} < \infty$.

In this paper, we generalize this result by proving that it is valid for any ℓ^r space, where $r \in [1, +\infty]$. This paper is organized as follows. In section 1, we recall basic concepts of Besov and Lizorkin-Triebel spaces, the decomposition of Littlewood-Paley, and some notations that will be needed throughout this paper. In section 2, we give a generalization of Bourdaud theorem of a localization property of Besov spaces on the ℓ^r space, where $r \in [1, +\infty]$. Also, we show that the Lizorkin-Triebel spaces are localizable in the ℓ^p norm. Finally, we present some conclusions and discuss future research in section 3.

1 PRELIMINARIES AND NOTATIONS

This section contains the basic definitions and notations that will be needed throughout this paper.

1.1 Notations

We note (e_1, \dots, e_n) the canonical basis of \mathbb{R}^n , $x \cdot y = x_1 y_1 + \dots + x_n y_n$ the scalar product in \mathbb{R}^n , and for $\alpha \in \mathbb{N}^n$, $|\alpha| = \alpha_1 + \dots + \alpha_n$ $\frac{\partial^{|\alpha|} f}{\partial^{|\alpha|} x_1 \dots \partial^{|\alpha|} x_n}$ the partial derivative of the function f is denoted by $\partial^\alpha f$.

If $f : \mathbb{R}^n \rightarrow \mathbb{C}$, the support of f denoted by $\text{supp } f$. $\mathcal{D}(\mathbb{R}^n)$ is the space of test functions, i.e., of smooth functions which have compact support, $\mathcal{D}'(\mathbb{R}^n)$ is the dual of $\mathcal{D}(\mathbb{R}^n)$. $\mathcal{S}(\mathbb{R}^n)$ is the Schwartz space of functions $\mathcal{C}^\infty(\mathbb{R}^n)$ rapidly decreasing on \mathbb{R}^n , the dual $\mathcal{S}'(\mathbb{R}^n)$ is the space of tempered distributions.

If $f \in \mathcal{S}(\mathbb{R}^n)$, then its Fourier transform defined by:

$$\mathcal{F}(f(x))(\xi) = \int_{\mathbb{R}^n} \exp(-ix \cdot \xi) f(x) dx$$

and its inverse Fourier transform defined by:

$$\mathcal{F}^{-1}(\widehat{f}(\xi))(x) = (2\pi)^{-n} \int_{\mathbb{R}^n} \exp(ix \cdot \xi) \widehat{f}(\xi) d\xi.$$

Let A_1 and A_2 two spaces, we say that $A_1 \hookrightarrow A_2$ if there exists $c > 0$ such that $\|\cdot\|_{A_2} \leq c \|\cdot\|_{A_1}$. p' is the conjugate exponent of p , $\frac{1}{p} + \frac{1}{p'} = 1$ where $p \in [1, +\infty]$.

Let $k \in \mathbb{Z}^n$, τ_k is the translation operator defined by $\tau_k f(\cdot) = f(\cdot - k)$. L^p is the space of the measurable functions f such that $\|f\|_{L^p} = (\int_{\mathbb{R}^n} |f(x)|^p dx)^{\frac{1}{p}} < \infty$. ℓ^q is the space of sequences $(a_k)_k$ such that $\|(a_k)\|_{\ell^q} = (\sum_{k=0}^{\infty} |a_k|^q)^{\frac{1}{q}} < \infty$.

Let $0 < p \leq \infty$, $0 < q \leq \infty$, so

$$\|f_k\|_{\ell^q(L^p)} = \left(\sum_{k=0}^{\infty} \|f_k(x)\|_p^q \right)^{\frac{1}{q}} < \infty, \quad \|f_k\|_{L^p(\ell^q)} = \left\| \left(\sum_{k=0}^{\infty} |f_k(x)|^q \right)^{\frac{1}{q}} \right\|_p < \infty.$$

1.2 The decomposition of Littlewood-Paley

Let $\varphi \in \mathcal{S}(\mathbb{R}^n)$, which satisfy the conditions:

- (i) $\text{supp } \varphi \subset \{\xi \in \mathbb{R}^n : 1 \leq |\xi| \leq 3\}$,
- (ii) $\varphi(\xi) > 0$, for $1 \leq |\xi| \leq 3$,

(iii) $\sum_{j \in \mathbb{Z}} \varphi(2^{-j}\xi) = 1$, for, $\xi \in \mathbb{R}^n \setminus \{0\}$.

The construction of φ does not pose any difficulty, see for example [3]. We put $\varphi(\xi) = 1 - \sum_{j=1}^{\infty} \varphi(2^{-j}\xi)$, then it follows that the function $\varphi \in C^\infty(\mathbb{R}^n)$, such that $\text{supp } \varphi \subset \{\xi \in \mathbb{R}^n : |\xi| \leq 3\}$. In the following, we fix the partition of the unit and we obtain:

$$\varphi(\xi) + \sum_{j=1}^{\infty} \varphi(2^{-j}\xi) = 1 \quad (\text{for all } \xi \in \mathbb{R}^n).$$

To this partition, we associate a sequence of convolution operators $\Delta_j : S' \rightarrow C^\infty$, defined by $\mathcal{F}(\Delta_j f)(\xi) = \varphi(2^{-j}\xi)\widehat{f}(\xi)$, for $j = 1, 2, \dots$ and $\mathcal{F}(\Delta_0 f)(\xi) = \varphi(\xi)\widehat{f}(\xi)$. Also, we define the operators Q_k by $\mathcal{F}(Q_k f)(\xi) = \varphi(2^{-k}\xi)\widehat{f}(\xi)$, for, $k = 1, 2, \dots$ for all $f \in S'$, the decomposition of f of the Littlewood-Paley type given by

$$f = \sum_{j \geq 0} \Delta_j f. \tag{1}$$

The series (1) converges in the sense of tempered distributions. The series (1) can be written as

$$f = Q_k f + \sum_{j \geq k+1} \Delta_j f,$$

This formula is valid for any $f \in S'$ and $k \in \mathbb{N}$, such that $Q_k f = \sum_{j \leq k} \Delta_j f$.

Definition 1 ([10]). Let $f \in S'$ and $a > 0$. We define the maximal operators associated to the Δ_k and Q_k by

$$\Delta_k^{*,a} f(x) = \sup_{y \in \mathbb{R}^n} \frac{|\Delta_k f(x-y)|}{(1+2^k|y|)^a} \quad \text{and} \quad Q_k^{*,a} f(x) = \sup_{y \in \mathbb{R}^n} \frac{|Q_k f(x-y)|}{(1+2^k|y|)^a}.$$

Notation: The symbol \hookrightarrow indicates that the embedding is continuous.

Definition 2 ([8]). Let $s \in \mathbb{R}, p, q \in [1, +\infty]$. The Besov space $B_{p,q}^s(\mathbb{R}^n)$ is the set of all $f \in S'(\mathbb{R}^n)$ satisfying

$$\|f\|_{B_{p,q}^s(\mathbb{R}^n)} = \begin{cases} (\sum_{j \geq 0} (2^{sj} \|\Delta_j f\|_p)^q)^{\frac{1}{q}} < +\infty, & \text{for } q \neq \infty, \\ \sup_{j \geq 0} (2^{sj} \|\Delta_j f\|_p) < +\infty, & \text{for } q = \infty. \end{cases} \tag{2}$$

Definition 3 ([8]). Let $s \in \mathbb{R}, p \in [1, +\infty[$ and $q \in [1, +\infty]$. The Lizorkin-Triebel space $F_{p,q}^s(\mathbb{R}^n)$ is the set of all $f \in S'(\mathbb{R}^n)$ satisfying

$$\|f\|_{F_{p,q}^s(\mathbb{R}^n)} = \begin{cases} \|(\sum_{j \geq 0} (2^{sj} |\Delta_j f|)^q)^{\frac{1}{q}}\|_p < +\infty, & \text{for } q \neq \infty. \\ \|\sup_{j \geq 0} (2^{sj} |\Delta_j f|)\|_p < +\infty, & \text{for } q = \infty. \end{cases} \tag{3}$$

Remark 1. In the formula (2)(resp. (3)), we can replace Δ_j by $\Delta_j^{*,a}$ with $a > \frac{n}{p}$ (resp. $a > \frac{n}{\min(p,q)}$), and we obtain an equivalent norm in $B_{p,q}^s(\mathbb{R}^n)$ (resp. $F_{p,q}^s(\mathbb{R}^n)$).

For more details, see Peetre [7] and Triebel [10].

Proposition 1 ([2]). Let $s \in \mathbb{R}$.

(i) for all $\gamma > 1$, there exists, $c > 0$, such that for any sequence of functions $(f_j)_{j \geq 0}$, where $\text{supp } \mathcal{F}f_j \subset \{\xi : \gamma^{-1}2^j \leq |\xi| \leq \gamma 2^j\}$, we have

$$\left\| \sum_{j=0}^{\infty} f_j \right\|_{B_{p,q}^s} \leq c \left(\sum_{j=0}^{\infty} 2^{sjq} \|f_j\|_p^q \right)^{\frac{1}{q}}.$$

(ii) for all $a > 1$, there exists, $c > 0$, such that for any sequence of functions $(f_j)_{j \in \mathbb{N}}$, where $\text{supp } \mathcal{F}f_j \subset \{\xi : a^{-1}2^j \leq |\xi| \leq a 2^j\}$, we have

$$\left\| \sum_{j=0}^{\infty} f_j \right\|_{F_{p,q}^s} \leq c \left\| \left(\sum_{j=0}^{\infty} 2^{sjq} |f_j|^q \right)^{\frac{1}{q}} \right\|_p.$$

2 LOCALIZATION OF BESOV SPACES

In this section, we give a generalization of Bourdaud theorem of a localization property of Besov spaces on the ℓ^r space, where $r \in [1, +\infty]$. Also, we show that the Lizorkin-Triebel spaces are localizable in the ℓ^p norm. We start with these important concepts.

Let E be a Banach space of distributions. We associate on the space E the following hypothesis.

- (1) Translation invariance; if we denote τ_x the operator given by $\tau_x f(t) = f(x - t)$, then τ_x is an isometric of E ;
- (2) Localization invariance; for all $f \in E$ and $\varphi \in \mathcal{D}(\mathbb{R}^n)$, we have that $\varphi f \in E$.

Let $\varphi \in \mathcal{D}(\mathbb{R}^n)$. The notion of localized is defined by $f_x = \tau_x \varphi \cdot f$, it follows immediately from the hypothesis 1 and 2 that the family $(f_x)_{x \in \mathbb{R}^n}$ is bounded in E . We consider the set A as the class of all the functions $\varphi \in \mathcal{D}(\mathbb{R}^n)$ satisfying

$$\sum_{k \in \mathbb{Z}^n} \varphi(x - k) = 1, \text{ for all } x \in \mathbb{R}^n.$$

Definition 4 ([1]). *Let E be a Banach space of distributions, E is localizable in the ℓ^p norm ($1 \leq p \leq \infty$), if there exists $\varphi \in A$ and a constant $c \geq 1$, such that*

$$c^{-1} \|f\|_E \leq \left(\sum_{k \in \mathbb{Z}^n} \|\tau_k \varphi \cdot f\|_E^p \right)^{\frac{1}{p}} \leq c \|f\|_E.$$

i.e., $E = (E)_{\ell^p}$, we denote by $(E)_{\ell^p}$ the distribution space of u such that

$$\|u\|_{(E)_{\ell^p}} = \left\| \left(\|\tau_k \varphi \cdot u\|_E \right)_{k \in \mathbb{Z}^n} \right\|_{\ell^p} < \infty.$$

Proposition 2 ([1]). *Let S be the Schwartz space, if the function $\theta \in S$ is not null on the support of φ , then we have:*

$$\|u\|_{(E)_{\ell^p}} \sim \left\| \left(\|\tau_k \theta \cdot u\|_E \right)_{k \in \mathbb{Z}^n} \right\|_{\ell^p}.$$

Proposition 3 ([1]). *Let $B_{p,q}^s$ be a Besov space, and N be a natural number fulfill $N > s$, and $\lambda, \mu \in S$, such that*

- (i) $\mu(\xi) \neq 0$, for $|\xi| \leq 3$,

(ii) $\lambda(\xi) \neq 0$, for $1 \leq |\xi| \leq 3$ and $\lambda^{(\alpha)}(0) = 0$ for $|\alpha| < N$.

We denote by $L_j (j \geq 1)$ the respective symbol operators $\lambda(2^{-j}\xi)$ and by L_0 the symbol operator $\mu(\xi)$, therefore

$$\|u\|_{B_{p,q}^s} \sim \|(2^{js} \|L_j u\|_p)_{j \in \mathbb{N}}\|_{\ell^q}.$$

In the following theorem we give a generalization of Bourdaud theorem of a localization property of Besov spaces on the ℓ^r spaces, by using Proposition 2 and Proposition 3.

Theorem 2. Let $p, q, r \in [1, +\infty]$, $s \in \mathbb{R}$, and $B_{p,q}^s$ and $(B_{p,q}^s)_{\ell^r}$ are respectively the Besov and localized Besov spaces. Then

- (i) $B_{p,q}^s \hookrightarrow (B_{p,q}^s)_{\ell^r}$, for $r \geq \max(p, q)$,
- (ii) $(B_{p,q}^s)_{\ell^r} \hookrightarrow B_{p,q}^s$ for $r \leq \min(p, q)$.

In particular, $B_{p,p}^s$ space is localizable in the ℓ^p norm.

Proof. (i) We will show that

$$\|u\|_{(B_{p,q}^s)_{\ell^r}} \leq c \|u\|_{B_{p,q}^s} \quad \text{for } c > 0.$$

By Proposition 1, it follows that

$$\left\| \sum_{j \geq 0} \tau_k \theta \cdot \Delta_j u \right\|_{B_{p,q}^s} \leq c \left(\sum_{j \geq 0} 2^{sjq} \|\tau_k \theta \cdot \Delta_j u\|_p^q \right)^{\frac{1}{q}}.$$

This implies that, $\|\tau_k \theta \cdot u\|_{B_{p,q}^s}^r \leq c \left(\sum_{j=0}^{\infty} 2^{sjq} \|\tau_k \theta \cdot \Delta_j u\|_p^q \right)^{\frac{r}{q}}$. Then it holds that

$$\left(\sum_{k \in \mathbb{Z}^n} \|\tau_k \theta \cdot u\|_{B_{p,q}^s}^r \right)^{\frac{1}{r}} \leq c \left(\sum_{k \in \mathbb{Z}^n} \left(\sum_{j=0}^{\infty} 2^{sjq} \|\tau_k \theta \cdot \Delta_j u\|_p^q \right)^{\frac{r}{q}} \right)^{\frac{1}{r}}.$$

Consequently

$$\left(\sum_{k \in \mathbb{Z}^n} \|\tau_k \theta \cdot u\|_{B_{p,q}^s}^r \right)^{\frac{1}{r}} \leq c \left(\|(2^{sj} \|\tau_k \theta \cdot \Delta_j u\|_p)_{k \in \mathbb{Z}^n}\|_{\ell^r(\ell^q)} \right). \quad (4)$$

Since, $r \geq \max(p, q)$ implies that $q \leq r$. Then from Minkowski inequality we have

$$\|(2^{sj} \|\tau_k \theta \cdot \Delta_j u\|_p)_{k \in \mathbb{Z}^n}\|_{\ell^r(\ell^q)} \leq c \|(2^{sj} \|\tau_k \theta \cdot \Delta_j u\|_p)_{k \in \mathbb{Z}^n}\|_{\ell^q(\ell^r)}.$$

So, we can see that the inequality (4) becomes as follows

$$\left(\sum_{k \in \mathbb{Z}^n} \|\tau_k \theta \cdot u\|_{B_{p,q}^s}^r \right)^{\frac{1}{r}} \leq c \left(\|(2^{sj} \|\tau_k \theta \cdot \Delta_j u\|_p)_{k \in \mathbb{Z}^n}\|_{\ell^q(\ell^r)} \right).$$

Consequently $\left(\sum_{k \in \mathbb{Z}^n} \|\tau_k \theta \cdot u\|_{B_{p,q}^s}^r \right)^{\frac{1}{r}} \leq c \left(\sum_{j=0}^{\infty} 2^{sjq} \left(\sum_{k \in \mathbb{Z}^n} \|\tau_k \theta \cdot \Delta_j u\|_p^q \right)^{\frac{1}{q}} \right)^{\frac{1}{q}}$. Therefore,

$$\left(\sum_{k \in \mathbb{Z}^n} \|\tau_k \theta \cdot u\|_{B_{p,q}^s}^r \right)^{\frac{1}{r}} \leq c \left(\sum_{j=0}^{\infty} 2^{sjq} (\|\tau_k \theta \cdot \Delta_j u\|_{\ell^r(L^p)}^q)^{\frac{1}{q}} \right). \quad (5)$$

Also, we have $r \geq \max(p, q)$ implies that $p \leq r$ i.e., $\ell^p \hookrightarrow \ell^r$, it follows that $\ell^p(L^p) \hookrightarrow \ell^r(L^p)$. Consequently $\|(\tau_k \theta \cdot \Delta_j u)_{k \in \mathbb{Z}^n}\|_{\ell^r(L^p)} \leq c \|(\tau_k \theta \cdot \Delta_j u)_{k \in \mathbb{Z}^n}\|_{\ell^p(L^p)}$. So, we can see that the inequality (5) becomes as follows $(\sum_{k \in \mathbb{Z}^n} \|\tau_k \theta \cdot u\|_{B_{p,q}^s}^r)^{\frac{1}{r}} \leq c (\sum_{j=0}^{\infty} 2^{sjq} (\|\tau_k \theta \cdot \Delta_j u\|_{\ell^p(L^p)}^q)^{\frac{1}{q}}$. Since L^p is a space localizable in the ℓ^p norm, then it holds that $\|\tau_k \theta \cdot \Delta_j u\|_{\ell^p(L^p)} \sim \|\Delta_j u\|_p$. Hence,

$$\left(\sum_{k \in \mathbb{Z}^n} \|\tau_k \theta \cdot u\|_{B_{p,q}^s}^r \right)^{\frac{1}{r}} \leq c \left(\sum_{j=0}^{\infty} 2^{sjq} \|\Delta_j u\|_p^q \right)^{\frac{1}{q}} \leq c \|u\|_{B_{p,q}^s}.$$

Thus, $B_{p,q}^s \hookrightarrow (B_{p,q}^s)_{\ell^r}$.

(ii) Now, we will show that

$$\|u\|_{B_{p,q}^s} \leq c \|u\|_{(B_{p,q}^s)_{\ell^r}}, \text{ for } c > 0.$$

Let $u \in \mathcal{S}'(\mathbb{R}^n)$. Then it holds that

$$\|L_j(u)\|_p = \|L_j(\sum_{k \in \mathbb{Z}^n} \tau_k \varphi \cdot u)\|_p = \left\| \sum_{k \in \mathbb{Z}^n} L_j(\tau_k \varphi \cdot u) \right\|_p \leq c \left(\sum_{k \in \mathbb{Z}^n} \|L_j(\tau_k \varphi \cdot u)\|_p^p \right)^{\frac{1}{p}}.$$

Since, $r \leq \min(p, q)$, it holds that $\ell^r \hookrightarrow \ell^p$ i.e.,

$$\|(\|L_j(\tau_k \varphi \cdot u)\|_p)_{k \in \mathbb{Z}^n}\|_{\ell^p} \leq c \|(\|L_j(\tau_k \varphi \cdot u)\|_p)_{k \in \mathbb{Z}^n}\|_{\ell^r}.$$

So, we have

$$\|L_j(u)\|_p \leq c \left(\sum_{k \in \mathbb{Z}^n} \|L_j(\tau_k \varphi \cdot u)\|_p^p \right)^{\frac{1}{p}} \leq c \left(\sum_{k \in \mathbb{Z}^n} \|L_j(\tau_k \varphi \cdot u)\|_p^r \right)^{\frac{1}{r}}.$$

This implies that

$$\left(\sum_{j=0}^{\infty} 2^{sjq} \|L_j u\|_p^q \right)^{\frac{1}{q}} \leq c \left(\sum_{j=0}^{\infty} 2^{sjq} \left(\sum_{k \in \mathbb{Z}^n} \|L_j(\tau_k \varphi \cdot u)\|_p^r \right)^{\frac{q}{r}} \right)^{\frac{1}{q}}.$$

Consequently

$$\left(\sum_{j=0}^{\infty} 2^{sjq} \|L_j u\|_p^q \right)^{\frac{1}{q}} \leq c \left(\| (2^{sj} \|L_j(\tau_k \varphi \cdot u)\|_p)_{k \in \mathbb{Z}^n} \|_{\ell^q(\ell^r)} \right). \quad (6)$$

Since, $r \leq \min(p, q)$, it holds that $r \leq q$, Then from Minkowski inequality we have

$$\| (2^{sj} \|L_j(\tau_k \varphi \cdot u)\|_p)_{k \in \mathbb{Z}^n} \|_{\ell^q(\ell^r)} \leq c \| (2^{sj} \|L_j(\tau_k \varphi \cdot u)\|_p)_{k \in \mathbb{Z}^n} \|_{\ell^r(\ell^q)}.$$

So, we can see that the inequality (6) becomes as follows

$$\left(\sum_{j=0}^{\infty} 2^{sjq} \|L_j u\|_p^q \right)^{\frac{1}{q}} \leq c \| (2^{sj} \|L_j(\tau_k \varphi \cdot u)\|_p)_{k \in \mathbb{Z}^n} \|_{\ell^r(\ell^q)}.$$

Consequently

$$\begin{aligned} \left(\sum_{j=0}^{\infty} 2^{sjq} \|L_j u\|_p^q \right)^{\frac{1}{q}} &\leq c \left(\sum_{k \in \mathbb{Z}^n} \left(\sum_{j=0}^{\infty} 2^{sjq} \|L_j(\tau_k \varphi \cdot u)\|_p^q \right)^{\frac{r}{q}} \right)^{\frac{1}{r}} \\ &\leq c \left(\sum_{k \in \mathbb{Z}^n} \|\tau_k \varphi \cdot u\|_{B_{p,q}^s}^r \right)^{\frac{1}{r}} \leq c \|u\|_{(B_{p,q}^s)_{\ell^r}}. \end{aligned}$$

Thus, $(B_{p,q}^s)_{\ell^r} \hookrightarrow B_{p,q}^s$. □

Remark 2. The generalization of Bourdaud Theorem given by Sickel and Smirnov in 1999 [9] using the wavelet method, the aim of this work is to generalize the same Theorem of a localization property by using a different method.

Theorem 3. Let $p, q \in [1, +\infty]$, $s \in \mathbb{R}$, and $F_{p,q}^s$ and $(F_{p,q}^s)_{\ell^p}$ are respectively the Lizorkin-Triebel and localized Lizorkin-Triebel spaces. Then the space $F_{p,q}^s$ is localizable in the ℓ^p norm. i.e., $F_{p,q}^s = (F_{p,q}^s)_{\ell^p}$.

Proof. (i) $(F_{p,q}^s)_{\ell^p} \hookrightarrow F_{p,q}^s$ We will show that

$$\|f\|_{F_{p,q}^s} \leq c \|f\|_{(F_{p,q}^s)_{\ell^p}}, \text{ for } c > 0.$$

From Definition 3, $\|f\|_{F_{p,q}^s} = \|(\sum_{j=0}^{\infty} 2^{sjq} |\Delta_j f|^q)^{\frac{1}{q}}\|_p$. We put $\Delta_j f = \sum_{k \in \mathbb{Z}^n} \tau_k \varphi \cdot \Delta_j f$, it follows that $\|f\|_{F_{p,q}^s} = \|(\sum_{j=0}^{\infty} (\sum_{k \in \mathbb{Z}^n} 2^{sj} |\tau_k \varphi \Delta_j f|^q)^{\frac{1}{q}})\|_p$. Consequently

$$\|f\|_{F_{p,q}^s} = \| \|2^{sj} (\tau_k \varphi \Delta_j f)_{k \in \mathbb{Z}^n}\|_{\ell^q(\ell^1)} \|_p.$$

Since, $1 \leq q$. Then from Minkowski inequality we have

$$\|f\|_{F_{p,q}^s} = \| \|2^{sj} (\tau_k \varphi \Delta_j f)_{k \in \mathbb{Z}^n}\|_{\ell^q(\ell^1)} \|_p \leq \| \|2^{sj} (\tau_k \varphi \Delta_j f)_{k \in \mathbb{Z}^n}\|_{\ell^1(\ell^q)} \|_p.$$

Consequently

$$\|f\|_{F_{p,q}^s} \leq c \| \sum_{k \in \mathbb{Z}^n} \sum_{j=0}^{\infty} 2^{sjq} |\tau_k \varphi \Delta_j f|^q \|^{\frac{1}{q}} \|_p \leq c (\sum_{k \in \mathbb{Z}^n} \| \sum_{j=0}^{\infty} 2^{sjq} |\tau_k \varphi \Delta_j f|^q \|^{\frac{1}{q}} \|_p)^{\frac{1}{p}}.$$

Hence, $\|f\|_{F_{p,q}^s} \leq c (\sum_{k \in \mathbb{Z}^n} \| \tau_k \varphi \cdot f \|_{F_{p,q}^s}^p)^{\frac{1}{p}}$. Thus, $(F_{p,q}^s)_{\ell^p} \hookrightarrow F_{p,q}^s$.

(ii) $F_{p,q}^s \hookrightarrow (F_{p,q}^s)_{\ell^p}$. Now, we will show that

$$\|f\|_{(F_{p,q}^s)_{\ell^p}} \leq c \|f\|_{F_{p,q}^s}, \text{ for } c > 0.$$

Let $p, q \in [1, +\infty]$ and $s \in \mathbb{R}$. Then it holds that

$$(\sum_{k \in \mathbb{Z}^n} \| \tau_k \varphi \cdot f \|_{F_{p,q}^s}^p)^{\frac{1}{p}} = (\sum_{k \in \mathbb{Z}^n} \| \tau_k \varphi \sum_{j=0}^{\infty} \Delta_j f \|_{F_{p,q}^s}^p)^{\frac{1}{p}} = (\sum_{k \in \mathbb{Z}^n} \| \sum_{j=0}^{\infty} \Delta_j f \tau_k \varphi \|_{F_{p,q}^s}^p)^{\frac{1}{p}}.$$

From Proposition 1, it follows that

$$\begin{aligned} (\sum_{k \in \mathbb{Z}^n} \| \tau_k \varphi \cdot f \|_{F_{p,q}^s}^p)^{\frac{1}{p}} &\leq c (\sum_{k \in \mathbb{Z}^n} \| (\sum_{j=0}^{\infty} 2^{sjq} |\Delta_j f \tau_k \varphi|^q)^{\frac{1}{q}} \|_p^p)^{\frac{1}{p}} \\ &\leq c (\sum_{k \in \mathbb{Z}^n} \| \tau_k \varphi (\sum_{j=0}^{\infty} 2^{sjq} |\Delta_j f|^q)^{\frac{1}{q}} \|_p^p)^{\frac{1}{p}}. \end{aligned}$$

Since L^p is a space localizable in the ℓ^p norm, then it holds that

$$(\sum_{k \in \mathbb{Z}^n} \| \tau_k \varphi \cdot f \|_{F_{p,q}^s}^p)^{\frac{1}{p}} \leq c \| (\sum_{j=0}^{\infty} 2^{sjq} |\Delta_j f|^q)^{\frac{1}{q}} \|_p \leq c \|f\|_{F_{p,q}^s}.$$

Thus, $F_{p,q}^s \hookrightarrow (F_{p,q}^s)_{\ell^p}$. □

3 CONCLUSION

In this work, we have generalized the Bourdaud theorem of a localization property of Besov spaces $B_{p,q}^s(\mathbb{R}^n)$ on the ℓ^r space, where $s \in \mathbb{R}$, $p, q, r \in [1, +\infty]$. Also, we have provided that the Lizorkin-Triebel spaces are localizable in the ℓ^p norm. In future work, we will investigate the localization property on other functional spaces.

REFERENCES

- [1] Bourdaud G. *Localisations des espaces de Besov*. Studia Math. 1988, **90** (2), 153–163. (in French)
- [2] Bourdaud G. *Analyse fonctionnelle dans l'espace Euclidien*. In: Publications mathématiques de l'Université Paris 7, 23. UER de mathématiques de Paris VII, Paris, 1979. (in French)
- [3] Bergh J., Löfstrom J. *Interpolation spaces*. In: Chenciner A., Coates J. Grundlehren der mathematischen Wissenschaften, 223. Springer, Heidelberg, 1976.
- [4] Dahlberg B.E.J. *A note on Sobolev spaces*. Proc. Sympos. Pure Math. 1979, **35** (1), 183–185.
- [5] Hazewinkel M. *Encyclopaedia of Mathematics*. Springer Science+Business Media, Dordrecht.
- [6] Marcus M., Mizel V.J. *Every superposition operator mapping one Sobolev space into another is continuous*. J. Funct. Anal. 1979, **33** (2), 217–229. doi: 10.1016/0022-1236(79)90113-7
- [7] Peetre J. *New thoughts on Besov spaces*. In: Duke University mathematics series, 1. Mathematics Department Duke University, Durham N.C., 1976.
- [8] Runst T., Sickel W. *Sobolev spaces of fractional order, Nemytzkij operators and nonlinear partial differential equations*. De Gruyter, Berlin, 1996.
- [9] Sickel W., Smirnov I. *Localization properties of Besov spaces and of its associated multiplier spaces*. Jenaer Schriften zur Mathematik und Informatik 1999, **Math/Inf/99/21**.
- [10] Triebel H. *Theory of Function Spaces*. Birkhäuser, Basel, 1983.

Received 13.02.2018

Revised 28.05.2018

Ферахтія Н., Аллауї С.Е. *Узагальнення локалізаційної властивості просторів Бесова* // Карпатські матем. публ. — 2018. — Т.10, №1. — С. 71–78.

Поняття локалізаційної властивості простору Бесова введено Г. Бурдо, введено таким чином, що простори Бесова $B_{p,q}^s(\mathbb{R}^n)$, де $s \in \mathbb{R}$ і $p, q \in [1, +\infty]$, такі, що $p \neq q$, є нелокалізованими у нормі ℓ^p . Пізніше він показав, що простори Бесова $B_{p,q}^s$ вкладені в локалізовані простори Бесова $(B_{p,q}^s)_{\ell^p}$ (тобто $B_{p,q}^s \hookrightarrow (B_{p,q}^s)_{\ell^p}$, при $p \geq q$). Також було показано, що локалізовані простори Бесова $(B_{p,q}^s)_{\ell^p}$ вкладені в простори Бесова $B_{p,q}^s$ (і.е., $(B_{p,q}^s)_{\ell^p} \hookrightarrow B_{p,q}^s$, при $p \leq q$). Зокрема $B_{p,p}^s$ є локалізованим в нормі ℓ^p , де ℓ^p простір послідовностей $(a_k)_k$ таких, що $\|(a_k)\|_{\ell^p} < \infty$. У цій статті ми узагальнили теорему Бурдо про локалізаційну властивість просторів Бесова $B_{p,q}^s(\mathbb{R}^n)$ на простір ℓ^r , де $r \in [1, +\infty]$. А точніше ми довели, що будь-який простір Бесова $B_{p,q}^s$ є вкладений в локалізований простір Бесова $(B_{p,q}^s)_{\ell^r}$ (і.е., $B_{p,q}^s \hookrightarrow (B_{p,q}^s)_{\ell^r}$, при $r \geq \max(p, q)$). Також ми показали, що будь-який локалізований простір Бесова $(B_{p,q}^s)_{\ell^r}$ вкладений в простір Бесова $B_{p,q}^s$ (тобто $(B_{p,q}^s)_{\ell^r} \hookrightarrow B_{p,q}^s$, при $r \leq \min(p, q)$). І на завершення було показано, що простори Лізоркіна-Трібела $F_{p,q}^s(\mathbb{R}^n)$, де $s \in \mathbb{R}$ і $p, q \in [1, +\infty]$ є локалізованими в нормі ℓ^p (тобто $F_{p,q}^s = (F_{p,q}^s)_{\ell^p}$).

Ключові слова і фрази: простори Бесова, простори Лізоркіна-Трібела, локалізаційна властивість.